

Conservative Solutions to a Nonlinear Variational Wave Equation

Alberto Bressan and Yuxi Zheng

Department of Mathematics, The Pennsylvania State University
E-mail: bressan@math.psu.edu; yzheng@math.psu.edu

January 25, 2005

Abstract

We establish the existence of a conservative weak solution to the Cauchy problem for the nonlinear variational wave equation $u_{tt} - c(u)(c(u)u_x)_x = 0$, for initial data of finite energy. Here $c(\cdot)$ is any smooth function with uniformly positive bounded values.

Mathematics Subject Classification (2000): 35Q35

Keywords: Existence, uniqueness, singularity, coordinate transformation

1 Introduction

We are interested in the Cauchy problem

$$u_{tt} - c(u)(c(u)u_x)_x = 0, \quad (1.1)$$

with initial data

$$u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x). \quad (1.2)$$

Throughout the following, we assume that $c : \mathbb{R} \mapsto \mathbb{R}_+$ is a smooth, bounded, uniformly positive function. Even for smooth initial data, it is well known that the solution can lose regularity in finite time ([12]). It is thus of interest to study whether the solution can be extended beyond the time when a singularity appears. This is indeed the main concern of the present paper. In ([5]) we considered the related equation

$$u_t + f(u)_x = \frac{1}{2} \int_0^x f''(u) u_x^2 dx \quad (1.3)$$

and constructed a semigroup of solutions, depending continuously on the initial data. Here we establish similar results for the nonlinear wave equation (1.1). By introducing

new sets of dependent and independent variables, we show that the solution to the Cauchy problem can be obtained as the fixed point of a contractive transformation. Our main result can be stated as follows.

Theorem 1. *Let $c : \mathbb{R} \mapsto [\kappa^{-1}, \kappa]$ be a smooth function, for some $\kappa > 1$. Assume that the initial data u_0 in (1.2) is absolutely continuous, and that $(u_0)_x \in \mathbf{L}^2$, $u_1 \in \mathbf{L}^2$. Then the Cauchy problem (1.1)-(1.2) admits a weak solution $u = u(t, x)$, defined for all $(t, x) \in \mathbb{R} \times \mathbb{R}$. In the t - x plane, the function u is locally Hölder continuous with exponent $1/2$. This solution $t \mapsto u(t, \cdot)$ is continuously differentiable as a map with values in $\mathbf{L}_{\text{loc}}^p$, for all $1 \leq p < 2$. Moreover, it is Lipschitz continuous w.r.t. the \mathbf{L}^2 distance, i.e.*

$$\|u(t, \cdot) - u(s, \cdot)\|_{\mathbf{L}^2} \leq L |t - s| \quad (1.4)$$

for all $t, s \in \mathbb{R}$. The equation (1.1) is satisfied in integral sense, i.e.

$$\iint \left[\phi_t u_t - (c(u)\phi)_x c(u) u_x \right] dx dt = 0 \quad (1.5)$$

for all test functions $\phi \in \mathcal{C}_c^1$. Concerning the initial conditions, the first equality in (1.2) is satisfied pointwise, while the second holds in $\mathbf{L}_{\text{loc}}^p$ for $p \in [1, 2[$.

Our constructive procedure yields solutions which depend continuously on the initial data. Moreover, the “energy”

$$\mathcal{E}(t) \doteq \frac{1}{2} \int \left[u_t^2(t, x) + c^2(u(t, x)) u_x^2(t, x) \right] dx \quad (1.6)$$

remains uniformly bounded. More precisely, one has

Theorem 2. *A family of weak solutions to the Cauchy problem (1.1)-(1.2) can be constructed with the following additional properties. For every $t \in \mathbb{R}$ one has*

$$\mathcal{E}(t) \leq \mathcal{E}_0 \doteq \frac{1}{2} \int \left[u_1^2(x) + c^2(u_0(x)) (u_0)_x^2(x) \right] dx. \quad (1.7)$$

Moreover, let a sequence of initial conditions satisfy

$$\|(u_0^n)_x - (u_0)_x\|_{\mathbf{L}^2} \rightarrow 0, \quad \|u_1^n - u_1\|_{\mathbf{L}^2} \rightarrow 0,$$

and $u_0^n \rightarrow u_0$ uniformly on compact sets, as $n \rightarrow \infty$. Then one has the convergence of the corresponding solutions $u^n \rightarrow u$, uniformly on bounded subsets of the t - x plane.

It appears in (1.7) that the total energy of our solutions may decrease in time. Yet, we emphasize that our solutions are *conservative*, in the following sense.

Theorem 3. *There exists a continuous family $\{\mu_t; t \in \mathbb{R}\}$ of positive Radon measures on the real line with the following properties.*

- (i) At every time t , one has $\mu_t(\mathbb{R}) = \mathcal{E}_0$.
- (ii) For each t , the absolutely continuous part of μ_t has density $\frac{1}{2}(u_t^2 + c^2 u_x^2)$ w.r.t. the Lebesgue measure.
- (iii) For almost every $t \in \mathbb{R}$, the singular part of μ_t is concentrated on the set where $c'(u) = 0$.

In other words, the total energy represented by the measure μ is conserved in time. Occasionally, some of this energy is concentrated on a set of measure zero. At the times τ when this happens, μ_τ has a non-trivial singular part and $\mathcal{E}(\tau) < \mathcal{E}_0$. The condition (iii) puts some restrictions on the set of such times τ . In particular, if $c'(u) \neq 0$ for all u , then this set has measure zero.

We point out that what we do is constructing a continuous semigroup of solutions. Uniqueness within a class of conservative solutions at this point is only a conjecture.

The paper is organized as follows. In the next two subsections we briefly discuss the physical motivations for the equation and recall some known results on its solutions. In Section 2 we introduce a new set of independent and dependent variables, and derive some identities valid for smooth solutions. We formulate a set of equations in the new variables which is equivalent to (1.1). In the new variables all singularities disappear: Smooth initial data lead to globally smooth solutions. In Section 3 we use a contractive transformation in a Banach space with a suitable weighted norm to show that there is a unique solution to the set of equations in the new variables, depending continuously on the data u_0, u_1 . Going back to the original variables u, t, x , remarkably, in Section 4 we establish the Hölder continuity of these solutions $u = u(t, x)$, and show that the integral equation (1.5) is satisfied. Moreover, in Section 5, we study the conservativeness of the solutions, establish the energy inequality and the Lipschitz continuity of the map $t \mapsto u(t, \cdot)$. This already yields a proof of Theorem 2. In Section 6 we study the continuity of the maps $t \mapsto u_x(t, \cdot)$, $t \mapsto u_t(t, \cdot)$, completing the proof of Theorem 1. The proof of Theorem 3 is given in Section 7.

1.1 Physical background of the equation

Equation (1.1) has several physical origins. In the context of nematic liquid crystals, it comes as follows. The mean orientation of the long molecules in a nematic liquid crystal is described by a director field of unit vectors, $\mathbf{n} \in \mathbb{S}^2$, the unit sphere. Associated with the director field \mathbf{n} , there is the well-known Oseen-Frank potential energy density W given by

$$W(\mathbf{n}, \nabla \mathbf{n}) = \alpha |\mathbf{n} \times (\nabla \times \mathbf{n})|^2 + \beta (\nabla \cdot \mathbf{n})^2 + \gamma (\mathbf{n} \cdot \nabla \times \mathbf{n})^2. \quad (1.8)$$

The positive constants α , β , and γ are elastic constants of the liquid crystal. For the special case $\alpha = \beta = \gamma$, the potential energy density reduces to

$$W(\mathbf{n}, \nabla \mathbf{n}) = \alpha |\nabla \mathbf{n}|^2,$$

which is the potential energy density used in harmonic maps into the sphere \mathbb{S}^2 . There are many studies on the constrained elliptic system of equations for \mathbf{n} derived through variational principles from the potential (1.8), and on the parabolic flow associated with it, see [3, 9, 10, 16, 22, 36] and references therein. In the regime in which inertia effects dominate viscosity, however, the propagation of the orientation waves in the director field may then be modeled by the least action principle (Saxton [29])

$$\frac{\delta}{\delta u} \int \{ \partial_t \mathbf{n} \cdot \partial_t \mathbf{n} - W(\mathbf{n}, \nabla \mathbf{n}) \} dx dt = 0, \quad \mathbf{n} \cdot \mathbf{n} = 1. \quad (1.9)$$

In the special case $\alpha = \beta = \gamma$, this variational principle (1.9) yields the equation for harmonic wave maps from (1+3)-dimensional Minkowski space into the two sphere, see [8, 31, 32] for example. For planar deformations depending on a single space variable x , the director field has the special form

$$\mathbf{n} = \cos u(x, t) \mathbf{e}_x + \sin u(x, t) \mathbf{e}_y,$$

where the dependent variable $u \in \mathbb{R}^1$ measures the angle of the director field to the x -direction, and \mathbf{e}_x and \mathbf{e}_y are the coordinate vectors in the x and y directions, respectively. In this case, the variational principle (1.9) reduces to (1.1) with the wave speed c given specifically by

$$c^2(u) = \alpha \cos^2 u + \beta \sin^2 u. \quad (1.10)$$

The equation (1.1) has interesting connections with long waves on a dipole chain in the continuum limit ([13], Zorski and Infeld [45], and Grundland and Infeld [14]), and in classical field theories and general relativity ([13]). We refer the interested reader to the article [13] for these connections.

This equation (1.1) compares interestingly with other well-known equations, e. g.

$$\partial_t^2 u - \partial_x [p(\partial_x u)] = 0, \quad (1.11)$$

where $p(\cdot)$ is a given function, considered by Lax [25], Klainerman and Majda [24], and Liu [28]. Second related equation is

$$\partial_t^2 u - c^2(u) \Delta u = 0 \quad (1.12)$$

considered by Lindblad [27], who established the global existence of smooth solutions of (1.12) with smooth, small, and spherically symmetric initial data in \mathbb{R}^3 , where the large-time decay of solutions in high space dimensions is crucial. The multi-dimensional generalization of equation (1.1),

$$\partial_t^2 u - c(u) \nabla \cdot (c(u) \nabla u) = 0, \quad (1.13)$$

contains a lower order term proportional to $cc' |\nabla u|^2$, which (1.12) lacks. This lower order term is responsible for the blow-up in the derivatives of u . Finally, we note that equation (1.1) also looks related to the perturbed wave equation

$$\partial_t^2 u - \Delta u + f(u, \nabla u, \nabla \nabla u) = 0, \quad (1.14)$$

where $f(u, \nabla u, \nabla \nabla u)$ satisfies an appropriate convexity condition (for example, $f = u^p$ or $f = a(\partial_t u)^2 + b|\nabla u|^2$) or some nullity condition. Blow-up for (1.14) with a convexity condition has been studied extensively, see [2, 11, 15, 20, 21, 26, 30, 33, 34] and Strauss [35] for more reference. Global existence and uniqueness of solutions to (1.14) with a nullity condition depend on the nullity structure and large time decay of solutions of the linear wave equation in higher dimensions (see Klainerman and Machedon [23] and references therein). Therefore (1.1) with the dependence of $c(u)$ on u and the possibility of sign changes in $c'(u)$ is familiar yet truly different.

Equation (1.1) has interesting asymptotic uni-directional wave equations. Hunter and Saxton ([17]) derived the asymptotic equations

$$(u_t + u^n u_x)_x = \frac{1}{2} n u^{n-1} (u_x)^2 \quad (1.15)$$

for (1.1) via weakly nonlinear geometric optics. We mention that the x -derivative of equation (1.15) appears in the high-frequency limit of the variational principle for the Camassa-Holm equation ([1, 6, 7]), which arises in the theory of shallow water waves. A construction of global solutions to the Camassa-Holm equations, based on a similar variable transformation as in the present paper, will appear in [4]

1.2 Known results

In [18], Hunter and Zheng established the global existence of weak solutions to (1.15) ($n = 1$) with initial data of bounded variations. It has also been shown that the dissipative solutions are limits of vanishing viscosity. Equation (1.15) ($n = 1$) is also shown to be completely integrable ([19]). In [37]–[44], Ping Zhang and Zheng study the global existence, uniqueness, and regularity of the weak solutions to (1.15) ($n = 1, 2$) with L^2 initial data, and special cases of (1.1). The study of the asymptotic equation has been very beneficial for both the blow-up result [12] and the current global existence result for the wave equation (1.1).

2 Variable Transformations

We start by deriving some identities valid for smooth solutions. Consider the variables

$$\begin{cases} R \doteq u_t + c(u)u_x, \\ S \doteq u_t - c(u)u_x, \end{cases} \quad (2.1)$$

so that

$$u_t = \frac{R + S}{2}, \quad u_x = \frac{R - S}{2c}. \quad (2.2)$$

By (1.1), the variables R, S satisfy

$$\begin{cases} R_t - cR_x = \frac{c'}{4c}(R^2 - S^2), \\ S_t + cS_x = \frac{c'}{4c}(S^2 - R^2). \end{cases} \quad (2.3)$$

Multiplying the first equation in (2.3) by R and the second one by S , we obtain balance laws for R^2 and S^2 , namely

$$\begin{cases} (R^2)_t - (cR^2)_x = \frac{c'}{2c}(R^2S - RS^2), \\ (S^2)_t + (cS^2)_x = -\frac{c'}{2c}(R^2S - RS^2). \end{cases} \quad (2.4)$$

As a consequence, the following quantities are conserved:

$$E \doteq \frac{1}{2}(u_t^2 + c^2u_x^2) = \frac{R^2 + S^2}{4}, \quad M \doteq -u_tu_x = \frac{S^2 - R^2}{4c}. \quad (2.5)$$

Indeed we have

$$\begin{cases} E_t + (c^2M)_x = 0, \\ M_t + E_x = 0. \end{cases} \quad (2.6)$$

One can think of $R^2/4$ as the energy density of backward moving waves, and $S^2/4$ as the energy density of forward moving waves.

We observe that, if R, S satisfy (2.3) and u satisfies (2.1b), then the quantity

$$F \doteq R - S - 2cu_x \quad (2.7)$$

provides solutions to the linear homogeneous equation

$$F_t - cF_x = \frac{c'}{2c}(R + S + 2cu_x)F. \quad (2.8)$$

In particular, if $F \equiv 0$ at time $t = 0$, the same holds for all $t > 0$. Similarly, if R, S satisfy (2.3) and u satisfies (2.1a), then the quantity

$$G \doteq R + S - 2u_t$$

provides solutions to the linear homogeneous equation

$$G_t + cG_x = \frac{c'}{2c}(R + S - 2cu_x)G.$$

In particular, if $G \equiv 0$ at time $t = 0$, the same holds for all $t > 0$. We thus have

Proposition 1. *Any smooth solution of (1.1) provides a solution to (2.1)–(2.3). Conversely, any smooth solution of (2.1b) and (2.3) (or (2.1a) and (2.3)) which satisfies (2.2b) (or (2.2a)) at time $t = 0$ provides a solution to (1.1).*

The main difficulty in the analysis of (1.1) is the possible breakdown of regularity of solutions. Indeed, even for smooth initial data, the quantities u_x, u_t can blow up in finite time. This is clear from the equations (2.3), where the right hand side grows quadratically, see ([12]) for handling change of signs of c' and interaction between R

and S . To deal with possibly unbounded values of R, S , it is convenient to introduce a new set of dependent variables:

$$w \doteq 2 \arctan R, \quad z \doteq 2 \arctan S,$$

so that

$$R = \tan \frac{w}{2}, \quad S = \tan \frac{z}{2}. \quad (2.9)$$

Using (2.3), we obtain the equations

$$w_t - c w_x = \frac{2}{1 + R^2} (R_t - c R_x) = \frac{c'}{2c} \frac{R^2 - S^2}{1 + R^2}, \quad (2.10)$$

$$z_t + c z_x = \frac{2}{1 + S^2} (S_t + c S_x) = \frac{c'}{2c} \frac{S^2 - R^2}{1 + S^2}. \quad (2.11)$$




Figure 1: Characteristic curves

To reduce the equation to a semilinear one, it is convenient to perform a further change of independent variables (fig. 1). Consider the equations for the forward and backward characteristics:

$$\dot{x}^+ = c(u), \quad \dot{x}^- = -c(u). \quad (2.12)$$

The characteristics passing through the point (t, x) will be denoted by

$$s \mapsto x^+(s, t, x), \quad s \mapsto x^-(s, t, x),$$

respectively. As coordinates (X, Y) of a point (t, x) we shall use the quantities

$$X \doteq \int_0^{x^-(0, t, x)} (1 + R^2(0, x)) dx, \quad Y \doteq \int_{x^+(0, t, x)}^0 (1 + S^2(0, x)) dx. \quad (2.13)$$

Of course this implies

$$X_t - c(u)X_x = 0, \quad Y_t + c(u)Y_x = 0, \quad (2.14)$$

$$(X_x)_t - (c X_x)_x = 0, \quad (Y_x)_t + (c Y_x)_x = 0. \quad (2.15)$$

Notice that

$$X_x(t, x) = \lim_{h \rightarrow 0} \frac{1}{h} \int_{x^-(0,t,x)}^{x^-(0,t,x+h)} (1 + R^2(0, x)) dx,$$

$$Y_x(t, x) = \lim_{h \rightarrow 0} \frac{1}{h} \int_{x^+(0,t,x)}^{x^+(0,t,x+h)} (1 + S^2(0, x)) dx.$$

For any smooth function f , using (2.14) one finds

$$\begin{aligned} f_t + cf_x &= f_X X_t + f_Y Y_t + cf_X X_x + cf_Y Y_x = (X_t + cX_x)f_X = 2cX_x f_X, \\ f_t - cf_x &= f_X X_t + f_Y Y_t - cf_X X_x - cf_Y Y_x = (Y_t - cY_x)f_Y = -2cY_x f_Y. \end{aligned} \quad (2.16)$$

We now introduce the further variables

$$p \doteq \frac{1 + R^2}{X_x}, \quad q \doteq \frac{1 + S^2}{-Y_x}. \quad (2.17)$$

Notice that the above definitions imply

$$(X_x)^{-1} = \frac{p}{1 + R^2} = p \cos^2 \frac{w}{2}, \quad (-Y_x)^{-1} = \frac{q}{1 + S^2} = q \cos^2 \frac{z}{2}. \quad (2.18)$$

From (2.10)-(2.11), using (2.16)-(2.18), we obtain

$$2c \frac{(1 + S^2)}{q} w_Y = \frac{c'}{2c} \frac{R^2 - S^2}{1 + R^2}, \quad 2c \frac{(1 + R^2)}{p} z_X = \frac{c'}{2c} \frac{S^2 - R^2}{1 + S^2}.$$

Therefore

$$\begin{cases} w_Y = \frac{c'}{4c^2} \frac{R^2 - S^2}{1 + R^2} \frac{q}{1 + S^2} = \frac{c'}{4c^2} \left(\sin^2 \frac{w}{2} \cos^2 \frac{z}{2} - \sin^2 \frac{z}{2} \cos^2 \frac{w}{2} \right) q, \\ z_X = \frac{c'}{4c^2} \frac{S^2 - R^2}{1 + S^2} \frac{p}{1 + R^2} = \frac{c'}{4c^2} \left(\sin^2 \frac{z}{2} \cos^2 \frac{w}{2} - \sin^2 \frac{w}{2} \cos^2 \frac{z}{2} \right) p. \end{cases} \quad (2.19)$$

Using trigonometric formulas, the above expressions can be further simplified as

$$\begin{cases} w_Y = \frac{c'}{8c^2} (\cos z - \cos w) q, \\ z_X = \frac{c'}{8c^2} (\cos w - \cos z) p. \end{cases}$$

Concerning the quantities p, q , we observe that

$$c_x = c' u_x = c' \frac{R - S}{2c}. \quad (2.20)$$

Using again (2.18) and (2.15) we compute

$$\begin{aligned}
p_t - c p_x &= (X_x)^{-1} 2R(R_t - cR_x) - (X_x)^{-2}[(X_x)_t - c(X_x)_x](1 + R^2) \\
&= (X_x)^{-1} 2R \frac{c'}{4c}(R^2 - S^2) - (X_x)^{-2}[c_x X_x](1 + R^2) \\
&= \frac{p}{1+R^2} 2R(R^2 - S^2) \frac{c'}{4c} - \frac{p}{1+R^2} \frac{c'}{2c}(R - S)(1 + R^2) \\
&= \frac{c'}{2c} \frac{p}{1+R^2} [S(1 + R^2) - R(1 + S^2)].
\end{aligned}$$

$$\begin{aligned}
q_t + c q_x &= (-Y_x)^{-1} 2S(S_t - cS_x) - (-Y_x)^{-2}[(-Y_x)_t + c(-Y_x)_x](1 + S^2) \\
&= (-Y_x)^{-1} 2S \frac{c'}{4c}(S^2 - R^2) - (-Y_x)^{-2}[c_x(Y_x)](1 + S^2) \\
&= \frac{q}{1+S^2} 2S(S^2 - R^2) \frac{c'}{4c} - \frac{q}{1+S^2} \frac{c'}{2c}(S - R)(1 + S^2) \\
&= \frac{c'}{2c} \frac{q}{1+S^2} [R(1 + S^2) - S(1 + R^2)].
\end{aligned}$$

In turn, this yields

$$\begin{aligned}
p_Y &= (p_t - c p_x) \frac{1}{2c(-Y_x)} = (p_t - c p_x) \frac{1}{2c} \frac{q}{1+S^2} \\
&= \frac{c'}{4c^2} \frac{S(1+R^2) - R(1+S^2)}{(1+R^2)(1+S^2)} pq = \frac{c'}{4c^2} \left[\frac{S}{1+S^2} - \frac{R}{1+R^2} \right] pq \\
&= \frac{c'}{4c^2} \left[\tan \frac{z}{2} \cos^2 \frac{z}{2} - \tan \frac{w}{2} \cos^2 \frac{w}{2} \right] pq = \frac{c'}{4c^2} \frac{\sin z - \sin w}{2} pq,
\end{aligned} \tag{2.21}$$

$$\begin{aligned}
q_X &= (q_t + c q_x) \frac{1}{2c(X_x)} = (q_t + c q_x) \frac{1}{2c} \frac{p}{1+R^2} \\
&= \frac{c'}{4c^2} \frac{R(1+S^2) - S(1+R^2)}{(1+S^2)(1+R^2)} pq = \frac{c'}{4c^2} \left[\frac{R}{1+R^2} - \frac{S}{1+S^2} \right] pq \\
&= \frac{c'}{4c^2} \left[\tan \frac{w}{2} \cos^2 \frac{w}{2} - \tan \frac{z}{2} \cos^2 \frac{z}{2} \right] pq = \frac{c'}{4c^2} \frac{\sin w - \sin z}{2} pq.
\end{aligned} \tag{2.22}$$

Finally, by (2.16) we have

$$\begin{aligned}
u_X &= (u_t + c u_x) \frac{1}{2c} \frac{p}{1+R^2} = \frac{1}{2c} \frac{R}{1+R^2} p = \frac{1}{2c} \left(\tan \frac{w}{2} \cos^2 \frac{w}{2} \right) p, \\
u_Y &= (u_t - c u_x) \frac{1}{2c} \frac{q}{1+S^2} = \frac{1}{2c} \frac{S}{1+S^2} q = \frac{1}{2c} \left(\tan \frac{z}{2} \cos^2 \frac{z}{2} \right) q.
\end{aligned} \tag{2.23}$$

Starting with the nonlinear equation (1.1), using X, Y as independent variables we thus obtain a semilinear hyperbolic system with smooth coefficients for the variables u, w, z, p, q . Using some trigonometric identities, the set of equations (2.19), (2.21)-(2.22) and (2.23) can be rewritten as

$$\begin{cases} w_Y = \frac{c'}{8c^2} (\cos z - \cos w) q, \\ z_X = \frac{c'}{8c^2} (\cos w - \cos z) p, \end{cases} \tag{2.24}$$

$$\begin{cases} p_Y = \frac{c'}{8c^2} [\sin z - \sin w] pq, \\ q_X = \frac{c'}{8c^2} [\sin w - \sin z] pq, \end{cases} \quad (2.25)$$

$$\begin{cases} u_X = \frac{\sin w}{4c} p, \\ u_Y = \frac{\sin z}{4c} q. \end{cases} \quad (2.26)$$

Remark 1. The function u can be determined by using either one of the equations in (2.26). One can easily check that the two equations are compatible, namely

$$\begin{aligned} u_{XY} &= -\frac{\sin w}{4c^2} c' u_Y p + \frac{\cos w}{4c} w_Y p + \frac{\sin w}{4c} p_Y \\ &= \frac{c'}{32c^3} [-2 \sin w \sin z + \cos w \cos z - \cos^2 w - \sin^2 w + \sin w \sin z] pq \\ &= \frac{c'}{32c^3} [\cos(w - z) - 1] pq \\ &= u_{YX}. \end{aligned} \quad (2.27)$$

Remark 2. We observe that the new system is invariant under translation by 2π in w and z . Actually, it would be more precise to work with the variables $w^\dagger \doteq e^{iw}$ and $z^\dagger \doteq e^{iz}$. However, for simplicity we shall use the variables w, z , keeping in mind that they range on the unit circle $[-\pi, \pi]$ with endpoints identified.

The system (2.24)-(2.26) must now be supplemented by non-characteristic boundary conditions, corresponding to (1.2). For this purpose, we observe that u_0, u_1 determine the initial values of the functions R, S at time $t = 0$. The line $t = 0$ corresponds to a curve γ in the (X, Y) plane, say

$$Y = \varphi(X), \quad X \in \mathbb{R},$$

where $Y \doteq \varphi(X)$ if and only if

$$X = \int_0^x (1 + R^2(0, x)) dx, \quad Y = - \int_0^x (1 + S^2(0, x)) dx \quad \text{for some } x \in \mathbb{R}.$$

We can use the variable x as a parameter along the curve γ . The assumptions $u_0 \in H^1$, $u_1 \in \mathbf{L}^2$ imply $R, S \in \mathbf{L}^2$; to fix the ideas, let

$$\mathcal{E}_0 \doteq \frac{1}{4} \int [R^2(0, x) + S^2(0, x)] dx < \infty. \quad (2.28)$$

The two functions

$$X(x) \doteq \int_0^x (1 + R^2(0, x)) dx, \quad Y(x) \doteq \int_x^0 (1 + S^2(0, x)) dx$$

are well defined and absolutely continuous. Clearly, X is strictly increasing while Y is strictly decreasing. Therefore, the map $X \mapsto \varphi(X)$ is continuous and strictly decreasing. From (2.28) it follows

$$|X + \varphi(X)| \leq 4\mathcal{E}_0. \quad (2.29)$$

As (t, x) ranges over the domain $[0, \infty[\times \mathbb{R}$, the corresponding variables (X, Y) range over the set

$$\Omega^+ \doteq \{(X, Y); Y \geq \varphi(X)\}. \quad (2.30)$$

Along the curve

$$\gamma \doteq \{(X, Y); Y = \varphi(X)\} \subset \mathbb{R}^2$$

parametrized by $x \mapsto (X(x), Y(x))$, we can thus assign the boundary data $(\bar{w}, \bar{z}, \bar{p}, \bar{q}, \bar{u}) \in \mathbf{L}^\infty$ defined by

$$\begin{cases} \bar{w} = 2 \arctan R(0, x), \\ \bar{z} = 2 \arctan S(0, x), \end{cases} \quad \begin{cases} \bar{p} \equiv 1, \\ \bar{q} \equiv 1, \end{cases} \quad \bar{u} = u_0(x). \quad (2.31)$$

We observe that the identity

$$F = \tan \frac{\bar{w}}{2} - \tan \frac{\bar{z}}{2} - 2c(\bar{u})\bar{u}_x = 0 \quad (2.32)$$

is identically satisfied along γ . A similar identity holds for G .

3 Construction of integral solutions

Aim of this section is to prove a global existence theorem for the system (2.24)-(2.26), describing the nonlinear wave equation in our transformed variables.

Theorem 4. *Let the assumptions in Theorem 1 hold. Then the corresponding problem (2.24)-(2.26) with boundary data (2.31) has a unique solution, defined for all $(X, Y) \in \mathbb{R}^2$.*

In the following, we shall construct the solution on the domain Ω^+ where $Y \geq \varphi(X)$. On the complementary set Ω^- where $Y < \varphi(X)$, the solution can be constructed in an entirely similar way.

Observing that all equations (2.24)-(2.26) have a locally Lipschitz continuous right hand side, the construction of a local solution as fixed point of a suitable integral transformation is straightforward. To make sure that this solution is actually defined on the whole domain Ω^+ , one must establish a priori bounds, showing that p, q remain bounded on bounded sets. This is not immediately obvious from the equations (2.25), because the right hand sides have quadratic growth.

The basic estimate can be derived as follows. Assume

$$C_0 \doteq \sup_{u \in \mathbb{R}} \left| \frac{c'(u)}{4c^2(u)} \right| < \infty. \quad (3.1)$$

From (2.25) it follows the identity

$$q_X + p_Y = 0.$$

In turn, this implies that the differential form $p dX - q dY$ has zero integral along every closed curve contained in Ω^+ . In particular, for every $(X, Y) \in \Omega^+$, consider the closed curve Σ (see fig. 2) consisting of:

- the vertical segment joining $(X, \varphi(X))$ with (X, Y) ,
- the horizontal segment joining (X, Y) with $(\varphi^{-1}(Y), Y)$
- the portion of boundary $\gamma = \{Y = \varphi(X)\}$ joining $(\varphi^{-1}(Y), Y)$ with $(X, \varphi(X))$.




Figure 2: The closed curve Σ

Integrating along Σ , recalling that $p = q = 1$ along γ and then using (2.29), we obtain

$$\begin{aligned} \int_{\varphi^{-1}(Y)}^X p(X', Y) dX' + \int_{\varphi(X)}^Y q(X, Y') dY' &= X - \varphi^{-1}(Y) + Y - \varphi(X) \\ &\leq 2(|X| + |Y| + 4\mathcal{E}_0). \end{aligned} \quad (3.2)$$

Using (3.1)-(3.2) in (2.25), since $p, q > 0$ we obtain the a priori bounds

$$\begin{aligned} p(X, Y) &= \exp \left\{ \int_{\varphi(X)}^Y \frac{c'(u)}{8c^2(u)} [\sin z - \sin w] q(X, Y') dY' \right\} \\ &\leq \exp \left\{ C_0 \int_{\varphi(X)}^Y q(X, Y') dY' \right\} \\ &\leq \exp \{ 2C_0(|X| + |Y| + 4\mathcal{E}_0) \}. \end{aligned} \quad (3.3)$$

Similarly,

$$q(X, Y) \leq \exp \{2C_0(|X| + |Y| + 4\mathcal{E}_0)\}. \quad (3.4)$$

Relying on (3.3)-(3.4), we now show that, on bounded sets in the X - Y plane, the solution of (2.24)–(2.26) with boundary conditions (2.31) can be obtained as the fixed point of a contractive transformation. For any given $r > 0$, consider the bounded domain

$$\Omega_r \doteq \{(X, Y); Y \geq \varphi(X), X \leq r, Y \leq r\}.$$

Introduce the space of functions

$$\Lambda_r \doteq \left\{ f : \Omega_r \mapsto \mathbb{R}; \|f\|_* \doteq \operatorname{ess\,sup}_{(X,Y) \in \Omega_r} e^{-\kappa(X+Y)} |f(X, Y)| < \infty \right\}$$

where κ is a suitably large constant, to be determined later. For $w, z, p, q, u \in \Lambda_r$, consider the transformation $\mathcal{T}(w, z, p, q, u) = (\tilde{w}, \tilde{z}, \tilde{p}, \tilde{q}, \tilde{u})$ defined by

$$\begin{cases} \tilde{w}(X, Y) = \bar{w}(X, \varphi(X)) + \int_{\varphi(X)}^Y \frac{c'(u)}{8c^2(u)} (\cos z - \cos w) q dY, \\ \tilde{z}(X, Y) = \bar{z}(\varphi^{-1}(Y), Y) + \int_{\varphi^{-1}(Y)}^X \frac{c'(u)}{8c^2(u)} (\cos w - \cos z) p dX, \end{cases} \quad (3.5)$$

$$\begin{cases} \tilde{p}(X, Y) = 1 + \int_{\varphi(X)}^Y \frac{c'(u)}{8c^2(u)} [\sin z - \sin w] \hat{p}\hat{q} dY, \\ \tilde{q}(X, Y) = 1 + \int_{\varphi^{-1}(Y)}^X \frac{c'(u)}{8c^2(u)} [\sin w - \sin z] \hat{p}\hat{q} dX, \end{cases} \quad (3.6)$$

$$\tilde{u}(X, Y) = \bar{u}(X, \varphi(X)) + \int_{\varphi(X)}^Y \frac{\sin z}{4c} q dY. \quad (3.7)$$

In (3.6), the quantities \hat{p}, \hat{q} are defined as

$$\hat{p} \doteq \min \{p, 2e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}\}, \quad \hat{q} \doteq \min \{q, 2e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}\}. \quad (3.8)$$

Notice that $\hat{p} = p, \hat{q} = q$ as long as the a priori estimates (3.3)-(3.4) are satisfied. Moreover, if in the equations (2.24)–(2.26) the variables p, q are replaced with \hat{p}, \hat{q} , then the right hand sides become uniformly Lipschitz continuous on bounded sets in the X - Y plane. A straightforward computation now shows that the map \mathcal{T} is a strict contraction on the space Λ_r , provided that the constant κ is chosen sufficiently big (depending on the function c and on r).

Obviously, if $r' > r$, then the solution of (3.5)–(3.7) on $\Omega_{r'}$ also provides the solution to the same equations on Ω_r , when restricted to this smaller domain. Letting $r \rightarrow \infty$, in the limit we thus obtain a unique solution (w, z, p, q, u) of (3.5)–(3.7), defined on the whole domain Ω^+ .

To prove that these functions satisfy the (2.24)–(2.26), we claim that $\hat{p} = p, \hat{q} = q$ at every point $(X, Y) \in \Omega^+$. The proof is by contradiction. If our claim does not hold, since the maps $Y \mapsto p(X, Y), X \mapsto q(X, Y)$ are continuous, we can find some point $(X^*, Y^*) \in \Omega^+$ such that

$$p(X, Y) \leq 2e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}, \quad q(X, Y) \leq 2e^{2C_0(|X|+|Y|+4\mathcal{E}_0)} \quad (3.9)$$

for all $(X, Y) \in \Omega^* \doteq \Omega^+ \cap \{X \leq X^*, Y \leq Y^*\}$, but either $p(X^*, Y^*) \geq \frac{3}{2}e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}$ or $q(X^*, Y^*) \geq \frac{3}{2}e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}$. By (3.9), we still have $\hat{p} = p$, $\hat{q} = q$ restricted to Ω^* , hence the equations (2.24)–(2.26) and the a priori bounds (3.3)–(3.4) remain valid. In particular, these imply

$$p(X^*, Y^*) \leq e^{2C_0(|X|+|Y|+4\mathcal{E}_0)}, \quad q(X^*, Y^*) \leq e^{2C_0(|X|+|Y|+4\mathcal{E}_0)},$$

reaching a contradiction. \square

Remark 3. In the solution constructed above, the variables w, z may well grow outside the initial range $] -\pi, \pi[$. This happens precisely when the quantities R, S become unbounded, i.e. when singularities arise.

For future reference, we state a useful consequence of the above construction.

Corollary 1. *If the initial data u_0, u_1 are smooth, then the solution (u, p, q, w, z) of (2.24)–(2.26), (2.31) is a smooth function of the variables (X, Y) . Moreover, assume that a sequence of smooth functions $(u_0^m, u_1^m)_{m \geq 1}$ satisfies*

$$u_0^m \rightarrow u_0, \quad (u_0^m)_x \rightarrow (u_0)_x, \quad u_1^m \rightarrow u_1$$

uniformly on compact subsets of \mathbb{R} . Then one has the convergence of the corresponding solutions:

$$(u^m, p^m, q^m, w^m, z^m) \rightarrow (u, p, q, w, z)$$

uniformly on bounded subsets of the X - Y plane.

We also remark that the equations (2.24)–(2.26) imply the conservation laws

$$q_X + p_Y = 0, \quad \left(\frac{q}{c}\right)_X - \left(\frac{p}{c}\right)_Y = 0. \quad (3.10)$$

4 Weak solutions, in the original variables

By expressing the solution $u(X, Y)$ in terms of the original variables (t, x) , we shall recover a solution of the Cauchy problem (1.1)–(1.2). This will provide a proof of Theorem 1.

As a preliminary, we examine the regularity of the solution (u, w, z, p, q) constructed in the previous section. Since the initial data $(u_0)_x$ and u_1 are only assumed to be in \mathbf{L}^2 , the functions w, z, p, q may well be discontinuous. More precisely, on bounded subsets of the X - Y plane, the equations (2.24)–(2.26) imply the following:

- The functions w, p are Lipschitz continuous w.r.t. Y , measurable w.r.t. X .
- The functions z, q are Lipschitz continuous w.r.t. X , measurable w.r.t. Y .
- The function u is Lipschitz continuous w.r.t. both X and Y .

The map $(X, Y) \mapsto (t, x)$ can be constructed as follows. Setting $f = x$, then $f = t$ in the two equations at (2.16), we find

$$\begin{cases} c = 2cX_x x_X, \\ -c = -2cY_x x_Y, \end{cases} \quad \begin{cases} 1 = 2cX_x t_X, \\ 1 = -2cY_x t_Y, \end{cases}$$

respectively. Therefore, using (2.18) we obtain

$$\begin{cases} x_X = \frac{1}{2X_x} = \frac{(1+\cos w)p}{4}, \\ x_Y = \frac{1}{2Y_x} = -\frac{(1+\cos z)q}{4}, \end{cases} \quad (4.1)$$

$$\begin{cases} t_X = \frac{1}{2cX_x} = \frac{(1+\cos w)p}{4c}, \\ t_Y = \frac{1}{-2cY_x} = \frac{(1+\cos z)q}{4c}. \end{cases} \quad (4.2)$$

For future reference, we write here the partial derivatives of the inverse mapping, valid at points where $w, z \neq -\pi$.

$$\begin{cases} X_x = \frac{2}{(1+\cos w)p}, \\ Y_x = -\frac{2}{(1+\cos z)q}, \end{cases} \quad \begin{cases} X_t = \frac{2c}{(1+\cos w)p}, \\ Y_t = \frac{2c}{(1+\cos z)q}. \end{cases} \quad (4.3)$$

We can now recover the functions $x = x(X, Y)$ by integrating one of the equations in (4.1). Moreover, we can compute $t = t(X, Y)$ by integrating one of the equations in (4.2). A straightforward calculation shows that the two equations in (4.1) are equivalent: differentiating the first w.r.t. Y or the second w.r.t. X one obtains the same expression.

$$\begin{aligned} x_{XY} &= \frac{(1+\cos w)p_Y}{4} - \frac{p \sin w w_Y}{4} \\ &= \frac{c' pq}{32c^2} [\sin z - \sin w + \sin(z - w)] = x_{YX}. \end{aligned}$$

Similarly, the equivalence of the two equations in (4.2) is checked by

$$\begin{aligned} t_{XY} - t_{YX} &= \left(\frac{x_X}{c}\right)_Y + \left(\frac{x_Y}{c}\right)_X = \frac{2}{c} x_{XY} - \left(\frac{x_X}{c^2} c' u_Y + \frac{x_Y}{c^2} c' u_X\right) \\ &= \frac{c' pq}{16c^3} [\sin z - \sin w + \sin(z - w)] \\ &\quad - \frac{c' pq}{16c^3} [(1 + \cos w) \sin z - (1 + \cos z) \sin w] = 0. \end{aligned}$$

In order to define u as a function of the original variables t, x , we should formally invert the map $(X, Y) \mapsto (t, x)$ and write $u(t, x) = u(X(t, x), Y(t, x))$. The fact that the above map may not be one-to-one does not cause any real difficulty. Indeed, given (t^*, x^*) , we can choose an arbitrary point (X^*, Y^*) such that $t(X^*, Y^*) = t^*$, $x(X^*, Y^*) = x^*$, and define $u(t^*, x^*) = u(X^*, Y^*)$. To prove that the values of u do not depend on the choice of (X^*, Y^*) , we proceed as follows. Assume that there are two

distinct points such that $t(X_1, Y_1) = t(X_2, Y_2) = t^*$, $x(X_1, Y_1) = x(X_2, Y_2) = x^*$. We consider two cases:

Case 1: $X_1 \leq X_2, Y_1 \leq Y_2$. Consider the set

$$\Gamma_{x^*} \doteq \left\{ (X, Y); \quad x(X, Y) \leq x^* \right\}$$

and call $\partial\Gamma_{x^*}$ its boundary. By (4.1), x is increasing with X and decreasing with Y . Hence, this boundary can be represented as the graph of a Lipschitz continuous function: $X - Y = \phi(X + Y)$. We now construct the Lipschitz continuous curve γ (fig. 3a) consisting of

- a horizontal segment joining (X_1, Y_1) with a point $A = (X_A, Y_A)$ on $\partial\Gamma_{x^*}$, with $Y_A = Y_1$,
- a portion of the boundary $\partial\Gamma_{x^*}$,
- a vertical segment joining (X_2, Y_2) to a point $B = (X_B, Y_B)$ on $\partial\Gamma_{x^*}$, with $X_B = X_2$.

We can obtain a Lipschitz continuous parametrization of the curve $\gamma : [\xi_1, \xi_2] \mapsto \mathbb{R}^2$ in terms of the parameter $\xi = X + Y$. Observe that the map $(X, Y) \mapsto (t, x)$ is constant along γ . By (4.1)-(4.2) this implies $(1 + \cos w)X_\xi = (1 + \cos z)Y_\xi = 0$, hence $\sin w \cdot X_\xi = \sin z \cdot Y_\xi = 0$. We now compute

$$\begin{aligned} u(X_2, Y_2) - u(X_1, Y_1) &= \int_\gamma (u_X dX + u_Y dY) \\ &= \int_{\xi_1}^{\xi_2} \left(\frac{p \sin w}{4c} X_\xi - \frac{q \sin z}{4c} Y_\xi \right) d\xi = 0, \end{aligned}$$

proving our claim.

Case 2: $X_1 \leq X_2, Y_1 \geq Y_2$. In this case, we consider the set

$$\Gamma_{t^*} \doteq \left\{ (X, Y); \quad t(X, Y) \leq t^* \right\},$$

and construct a curve γ connecting (X_1, Y_1) with (X_2, Y_2) as in fig. 3b. Details are entirely similar to Case 1.

We now prove that the function $u(t, x) = u(X(t, x), Y(t, x))$ thus obtained is Hölder continuous on bounded sets. Toward this goal, consider any characteristic curve, say $t \mapsto x^+(t)$, with $\dot{x}^+ = c(u)$. By construction, this is parametrized by the function $X \mapsto (t(X, \bar{Y}), x(X, \bar{Y}))$, for some fixed \bar{Y} . Recalling (2.16), (2.14), (2.18) and (2.26), we compute

$$\begin{aligned} \int_0^\tau [u_t + c(u)u_x]^2 dt &= \int_{X_0}^{X_\tau} (2cX_x u_X)^2 (2X_t)^{-1} dX \\ &= \int_{X_0}^{X_\tau} 2c \left(p \cos^2 \frac{w}{2} \right)^{-1} \left(\frac{p}{4c} 2 \sin \frac{w}{2} \cos \frac{w}{2} \right)^2 dX \quad (4.4) \\ &\leq \int_{X_0}^{X_\tau} \frac{p}{2c} dX \leq C_\tau, \end{aligned}$$




Figure 3: Paths of integration

for some constant C_τ depending only on τ . Similarly, integrating along any backward characteristics $t \mapsto x^-(t)$ we obtain

$$\int_0^\tau [u_t - c(u)u_x]^2 dt \leq C_\tau. \quad (4.5)$$

Since the speed of characteristics is $\pm c(u)$, and $c(u)$ is uniformly positive and bounded, the bounds (4.4)-(4.5) imply that the function $u = u(t, x)$ is Hölder continuous with exponent $1/2$. In turn, this implies that all characteristic curves are \mathcal{C}^1 with Hölder continuous derivative. Still from (4.4)-(4.5) it follows that the functions R, S at (2.1) are square integrable on bounded subsets of the t - x plane.

Finally, we prove that the function u provides a weak solution to the nonlinear wave equation (1.1). According to (1.5), we need to show that

$$\begin{aligned} 0 &= \iint \phi_t [(u_t + cu_x) + (u_t - cu_x)] - (c(u)\phi)_x [(u_t + cu_x) - (u_t - cu_x)] dxdt \\ &= \iint [\phi_t - (c\phi)_x] (u_t + cu_x) dxdt + \iint [\phi_t + (c\phi)_x] (u_t - cu_x) dxdt \\ &= \iint [\phi_t - (c\phi)_x] R dxdt + \iint [\phi_t + (c\phi)_x] S dxdt. \end{aligned} \quad (4.6)$$

By (2.16), this is equivalent to

$$\iint \left\{ -2cY_x\phi_Y R + 2cX_x\phi_X S + c'(u_X X_x + u_Y Y_x)\phi(S - R) \right\} dxdt = 0. \quad (4.7)$$

It will be convenient to express the double integral in (4.7) in terms of the variables X, Y . We notice that, by (2.18) and (2.14),

$$dx dt = \frac{pq}{2c(1+R^2)(1+S^2)} dXdY.$$

Using (2.26) and the identities

$$\begin{cases} \frac{1}{1+R^2} = \cos^2 \frac{w}{2} = \frac{1+\cos w}{2}, & \frac{R}{1+R^2} = \frac{\sin w}{2}, \\ \frac{1}{1+S^2} = \cos^2 \frac{z}{2} = \frac{1+\cos z}{2}, & \frac{S}{1+S^2} = \frac{\sin z}{2}, \end{cases} \quad (4.8)$$

the double integral in (4.6) can thus be written as

$$\begin{aligned}
& \iint \left\{ 2c \frac{1+S^2}{q} \phi_Y R + 2c \frac{1+R^2}{p} \phi_X S + c' \left(\frac{\sin w}{4c} p \frac{1+R^2}{p} - \frac{\sin z}{4c} q \frac{1+S^2}{q} \right) \phi (S - R) \right\} \\
& \quad \cdot \frac{pq}{2c(1+R^2)(1+S^2)} dXdY \\
& = \iint \left\{ \frac{R}{1+R^2} p \phi_Y + \frac{S}{1+S^2} q \phi_X + \frac{c'pq}{8c^2} \left(\frac{\sin w}{1+S^2} - \frac{\sin z}{1+R^2} \right) \phi (S - R) \right\} dXdY \\
& = \iint \left\{ \frac{p \sin w}{2} \phi_Y + \frac{q \sin z}{2} \phi_X \right. \\
& \quad \left. + \frac{c'pq}{8c^2} \left(\sin w \sin z - \sin w \cos^2 \frac{z}{2} \tan \frac{w}{2} - \sin z \cos^2 \frac{w}{2} \tan \frac{z}{2} \right) \phi \right\} dXdY \\
& = \iint \left\{ \frac{p \sin w}{2} \phi_Y + \frac{q \sin z}{2} \phi_X + \frac{c'pq}{8c^2} [\cos(w+z) - 1] \phi \right\} dXdY.
\end{aligned} \tag{4.9}$$

Recalling (2.30), one finds

$$\begin{aligned}
\left(\frac{p \sin w}{2} \right)_Y + \left(\frac{q \sin z}{2} \right)_X & = (2c u_X)_Y + (2c u_Y)_X \\
& = 4c' u_X u_Y + 4c u_{XY} \\
& = \frac{c'pq}{4c^2} \sin w \sin z + \frac{c'pq}{8c^2} [\cos(w-z) - 1] \\
& = \frac{c'pq}{8c^2} [\cos(w+z) - 1].
\end{aligned} \tag{4.10}$$

Together, (4.9) and (4.10) imply (4.7) and hence (4.6). This establishes the integral equation (1.5) for every test function $\phi \in \mathcal{C}_c^1$.

5 Conserved quantities

From the conservation laws (3.10) it follows that the 1-forms $p dX - q dY$ and $\frac{p}{c} dX + \frac{q}{c} dY$ are closed, hence their integrals along any closed curve in the X - Y plane vanish. From the conservation laws at (2.6), it follows that the 1-forms

$$E dx - (c^2 M) dt, \quad M dx - E dt \tag{5.1}$$

are also closed. There is a simple correspondence. In fact

$$E dx - (c^2 M) dt = \frac{p}{4} dX - \frac{q}{4} dY - \frac{1}{2} dx \quad - M dx + E dt = \frac{p}{4c} dX + \frac{q}{4c} dY - \frac{1}{2} dt.$$

Recalling (4.1)-(4.2), these can be written in terms of the X - Y coordinates as

$$\frac{(1 - \cos w) p}{8} dX - \frac{(1 - \cos z) q}{8} dY, \tag{5.2}$$

$$\frac{(1 - \cos w) p}{8c} dX + \frac{(1 - \cos z) q}{8c} dY, \tag{5.3}$$

respectively. Using (2.24)–(2.26), one easily checks that these forms are indeed closed:

$$\left(\frac{(1 - \cos w)p}{8}\right)_Y = \frac{c'pq}{64c^2} [\sin z(1 - \cos w) - \sin w(1 - \cos z)] = -\left(\frac{(1 - \cos z)q}{8}\right)_X, \quad (5.4)$$

$$\left(\frac{(1 - \cos w)p}{8c}\right)_Y = \frac{c'pq}{64c^3} [\sin(w + z) - (\sin w + \sin z)] = \left(\frac{(1 - \cos z)q}{8c}\right)_X.$$

In addition, we have the 1-forms

$$dx = \frac{(1 + \cos w)p}{4} dX - \frac{(1 + \cos z)q}{4} dY, \quad (5.5)$$

$$dt = \frac{(1 + \cos w)p}{4c} dX + \frac{(1 + \cos z)q}{4c} dY, \quad (5.6)$$

which are obviously closed.




Figure 4:

The solutions $u = u(X, Y)$ constructed in Section 3 are *conservative*, in the sense that the integral of the form (5.2) along every Lipschitz continuous, closed curve in the X - Y plane is zero.

To prove the inequality (1.7), fix any $\tau > 0$. The case $\tau < 0$ is identical. For a given $r > 0$ arbitrarily large, define the set (fig. 4)

$$\Gamma \doteq \left\{ (X, Y); 0 \leq t(X, Y) \leq \tau, \quad X \leq r, \quad Y \leq r \right\} \quad (5.7)$$

By construction, the the map $(X, Y) \mapsto (t, x)$ will act as follows:

$$A \mapsto (\tau, a), \quad B \mapsto (\tau, b), \quad C \mapsto (0, c), \quad D \mapsto (0, d),$$

for some $a < b$ and $d < c$. Integrating the 1-form (5.2) along the boundary of Γ we obtain

$$\begin{aligned}
& \int_{AB} \frac{(1-\cos w)p}{8} dX - \frac{(1-\cos z)q}{8} dY \\
&= \int_{DC} \frac{(1-\cos w)p}{8} dX - \frac{(1-\cos z)q}{8} dY - \int_{DA} \frac{(1-\cos w)p}{8} dX - \int_{CB} \frac{(1-\cos z)q}{8} dY \\
&\leq \int_{DC} \frac{(1-\cos w)p}{8} dX - \frac{(1-\cos z)q}{8} dY \\
&= \int_d^c \frac{1}{2} \left[u_t^2(0, x) + c^2(u(0, x)) u_x^2(0, x) \right] dx.
\end{aligned} \tag{5.8}$$

On the other hand, using (5.5) we compute

$$\begin{aligned}
& \int_a^b \frac{1}{2} \left[u_t^2(\tau, x) + c^2(u(\tau, x)) u_x^2(\tau, x) \right] dx \\
&= \int_{AB \cap \{\cos w \neq -1\}} \frac{(1-\cos w)p}{8} dX + \int_{AB \cap \{\cos z \neq -1\}} \frac{(1-\cos z)q}{8} dY \\
&\leq \mathcal{E}_0.
\end{aligned} \tag{5.9}$$

Notice that the last relation in (5.8) is satisfied as an equality, because at time $t = 0$, along the curve γ_0 the variables w, z never assume the value $-\pi$. Letting $r \rightarrow +\infty$ in (5.7), one has $a \rightarrow -\infty, b \rightarrow +\infty$. Therefore (5.8) and (5.9) together imply $\mathcal{E}(t) \leq \mathcal{E}_0$, proving (1.7).




Figure 5: Proving Lipschitz continuity

We now prove the Lipschitz continuity of the map $t \mapsto u(t, \cdot)$ in the \mathbf{L}^2 distance. For this purpose, for any fixed time τ , we let $\mu_\tau = \mu_\tau^- + \mu_\tau^+$ be the positive measure on the real line defined as follows. In the smooth case,

$$\mu_\tau^-(]a, b[) = \frac{1}{4} \int_a^b R^2(\tau, x) dx, \quad \mu_\tau^+(]a, b[) = \frac{1}{4} \int_a^b S^2(\tau, x) dx. \tag{5.10}$$

To define μ_τ^\pm in the general case, let γ_τ be the boundary of the set

$$\Gamma_\tau \doteq \{(X, Y); t(X, Y) \leq \tau\}. \tag{5.11}$$

Given any open interval $]a, b[$, let $A = (X_A, Y_A)$ and $B = (X_B, Y_B)$ be the points on γ_τ such that

$$\begin{aligned} x(A) = a, & \quad X_P - Y_P \leq X_A - Y_A \quad \text{for every point } P \in \gamma_\tau \text{ with } x(P) \leq a, \\ x(B) = b, & \quad X_P - Y_P \geq X_B - Y_B \quad \text{for every point } P \in \gamma_\tau \text{ with } x(P) \geq b. \end{aligned}$$

Then

$$\mu_\tau(]a, b[) = \mu_\tau^-(]a, b[) + \mu_\tau^+(]a, b[), \quad (5.12)$$

where

$$\mu_\tau^-(]a, b[) \doteq \int_{AB} \frac{(1 - \cos w)p}{8} dX \quad \mu_\tau^+(]a, b[) \doteq - \int_{AB} \frac{(1 - \cos z)q}{8} dY. \quad (5.13)$$

Recalling the discussion at (5.1)–(5.2), it is clear that μ_τ^- , μ_τ^+ are bounded, positive measures, and $\mu_\tau(\mathbb{R}) = \mathcal{E}_0$, for all τ . Moreover, by (5.10) and (2.5),

$$\int_a^b c^2 u_x^2 dx \leq \int_a^b \frac{1}{2} (R^2 + S^2) dx \leq 2\mu(]a, b[).$$

For any $a < b$, this yields the estimate

$$|u(\tau, b) - u(\tau, a)|^2 \leq |b - a| \int_a^b u_x^2(\tau, y) dy \leq 2\kappa^2 |b - a| \mu_\tau(]a, b[). \quad (5.14)$$

Next, for a given $h > 0$, $y \in \mathbb{R}$, we seek an estimate on the distance $|u(\tau+h, y) - u(\tau, y)|$. As in fig. 5, let $\gamma_{\tau+h}$ be the boundary of the set $\Gamma_{\tau+h}$, as in (5.11). Let $P = (P_X, P_Y)$ be the point on γ_τ such that

$$x(P) = y, \quad X_{P'} - Y_{P'} \leq X_P - Y_P \quad \text{for every point } P' \in \gamma_\tau \text{ with } x(P') \leq x,$$

Similarly, let $Q = (Q_X, Q_Y)$ be the point on $\gamma_{\tau+h}$ such that

$$x(Q) = y, \quad X_{Q'} - Y_{Q'} \leq X_Q - Y_Q \quad \text{for every point } Q' \in \gamma_{\tau+h} \text{ with } x(Q') \leq y.$$

Notice that $X_P \leq X_Q$ and $Y_P \leq Y_Q$. Let $P^+ = (X^+, Y^+)$ be a point on γ_τ with $X^+ = X_Q$, and let $P^- = (X^-, Y^-)$ be a point on γ_τ with $Y^- = Y_Q$. Notice that $x(P^+) \in]y, y + \kappa h[$, because the point $(\tau, x(Q))$ lies on some characteristic curve with speed $-c(u) > -\kappa$, passing through the point $(\tau+h, y)$. Similarly, $x(P^-) \in]y - \kappa h, y[$. Recalling that the forms in (5.2) and (5.6) are closed, we obtain the estimate

$$\begin{aligned} |u(Q) - u(P^+)| &\leq \int_{Y^+}^{Y_Q} |u_Y(X_Q, Y)| dY \\ &= \int_{Y^+}^{Y_Q} \left| \frac{\sin z}{4c} q \right| dY \\ &= \int_{Y^+}^{Y_Q} \left(\frac{(1+\cos z)q}{4c} \right)^{1/2} \left(\frac{(1-\cos z)q}{4} \right)^{1/2} dY \\ &\leq \left(\int_{Y^+}^{Y_Q} \frac{(1+\cos z)q}{4c} dY \right)^{1/2} \cdot \left(\int_{Y^+}^{Y_Q} \frac{(1-\cos z)q}{4} dY \right)^{1/2} \\ &\leq \left\{ \int_{P^- P^+} \left[\frac{(1-\cos w)p}{4} dX - \frac{(1-\cos z)q}{4} dY \right] \right\}^{1/2} \cdot h^{1/2}. \end{aligned} \quad (5.15)$$

The last term in (5.15) contains the integral of the 1-form at (5.2), along the curve γ_τ , between P^- and P^+ . Recalling the definition (5.12)–(5.13) and the estimate (5.14), we obtain the bound

$$\begin{aligned} |u(\tau + h, x) - u(\tau, x)|^2 &\leq 2|u(Q) - u(P^+)|^2 + 2|u(P^+) - u(P)|^2 \\ &\leq 4h \cdot \mu_\tau([x - \kappa h, x + \kappa h]) + 4\kappa^2 \cdot (\kappa h) \cdot \mu_\tau([x, x + h]). \end{aligned} \quad (5.16)$$

Therefore, for any $h > 0$,

$$\begin{aligned} \|u(\tau + h, \cdot) - u(\tau, \cdot)\|_{\mathbf{L}^2} &= \left(\int |u(\tau + h, x) - u(\tau, x)|^2 dx \right)^{1/2} \\ &\leq \left(\int 4(1 + \kappa^3)h \cdot \mu_\tau([x - \kappa h, x + \kappa h]) dx \right)^{1/2} \\ &= \left(4(\kappa^3 + 1)h^2 \mu_\tau(\mathbb{R}) \right)^{1/2} \\ &= h \cdot [4(\kappa^3 + 1) \mathcal{E}_0]^{1/2}. \end{aligned} \quad (5.17)$$

This proves the uniform Lipschitz continuity of the map $t \mapsto u(t, \cdot)$, stated at (1.4).

6 Regularity of trajectories

In this section we prove the continuity of the functions $t \mapsto u_t(t, \cdot)$ and $t \mapsto u_x(t, \cdot)$, as functions with values in \mathbf{L}^p . This will complete the proof of Theorem 1.

We first consider the case where the initial data $(u_0)_x, u_1$ are smooth with compact support. In this case, the solution $u = u(X, Y)$ remains smooth on the entire X - Y plane. Fix a time τ and let γ_τ be the boundary of the set Γ_τ , as in (5.11). We claim that

$$\left. \frac{d}{dt} u(t, \cdot) \right|_{t=\tau} = u_t(\tau, \cdot) \quad (6.1)$$

where, by (2.14), (2.18) and (2.26),

$$\begin{aligned} u_t(\tau, x) &\doteq u_X X_t + u_Y Y_t \\ &= \frac{p \sin w}{4c} \frac{2c}{p(1+\cos w)} + \frac{q \sin z}{4c} \frac{2c}{q(1+\cos z)} = \frac{\sin w}{2(1+\cos w)} + \frac{\sin z}{2(1+\cos z)}. \end{aligned} \quad (6.2)$$

Notice that (6.2) defines the values of $u_t(\tau, \cdot)$ at almost every point $x \in \mathbb{R}$, i.e. at all points outside the support of the singular part of the measure μ_τ defined at (5.12). By the inequality (1.7), recalling that $c(u) \geq \kappa^{-1}$, we obtain

$$\int_{\mathbb{R}} |u_t(\tau, x)|^2 dx \leq \kappa^2 \mathcal{E}(\tau) \leq \kappa^2 \mathcal{E}_0. \quad (6.3)$$

To prove (6.1), let any $\varepsilon > 0$ be given. There exist finitely many disjoint intervals $[a_i, b_i] \subset \mathbb{R}$, $i = 1, \dots, N$, with the following property. Call A_i, B_i the points on γ_τ such that $x(A_i) = a_i$, $x(B_i) = b_i$. Then one has

$$\min \{1 + \cos w(P), 1 + \cos z(P)\} < 2\varepsilon \quad (6.4)$$

at every point P on γ_τ contained in one of the arcs $A_i B_i$, while

$$1 + \cos w(P) > \varepsilon, \quad 1 + \cos z(P) > \varepsilon, \quad (6.5)$$

for every point P along γ_τ , not contained in any of the arcs $A_i B_i$. Call $J \doteq \cup_{1 \leq i \leq N} [a_i, b_i]$, $J' = \mathbb{R} \setminus J$, and notice that, as a function of the original variables, $u = u(t, x)$ is smooth in a neighborhood of the set $\{\tau\} \times J'$. Using Minkowski's inequality and the differentiability of u on J' , we can write

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{1}{h} \left(\int_{\mathbb{R}} \left| u(\tau + h, x) - u(\tau, x) - h u_t(\tau, x) \right|^p dx \right)^{1/p} \\ & \leq \lim_{h \rightarrow 0} \frac{1}{h} \left(\int_J \left| u(\tau + h, x) - u(\tau, x) \right|^p dx \right)^{1/p} + \left(\int_J |u_t(\tau, x)|^p dx \right)^{1/p} \end{aligned} \quad (6.6)$$

We now provide an estimate on the measure of the ‘‘bad’’ set J :

$$\begin{aligned} \text{meas}(J) &= \int_J dx = \sum_i \int_{A_i B_i} \frac{(1 + \cos w)^p}{4} dX - \frac{(1 + \cos z)^q}{4} dY \\ &\leq 2\varepsilon \sum_i \int_{A_i B_i} \frac{(1 - \cos w)^p}{4} dX - \frac{(1 - \cos z)^q}{4} dY \\ &\leq 2\varepsilon \int_{\gamma_\tau} \frac{(1 - \cos w)^p}{4} dX - \frac{(1 - \cos z)^q}{4} dY \leq 2\varepsilon \mathcal{E}_0. \end{aligned} \quad (6.7)$$

Now choose $q = 2/(2 - p)$ so that $\frac{p}{2} + \frac{1}{q} = 1$. Using Hölder's inequality with conjugate exponents $2/p$ and q , and recalling (5.17), we obtain

$$\begin{aligned} \int_J \left| u(\tau + h, x) - u(\tau, x) \right|^p dx &\leq \text{meas}(J)^{1/q} \cdot \left(\int_J \left| u(\tau + h, x) - u(\tau, x) \right|^2 dx \right)^{p/2} \\ &\leq [2\varepsilon \mathcal{E}_0]^{1/q} \cdot \left(\|u(\tau + h, \cdot) - u(\tau, \cdot)\|_{\mathbf{L}^2}^2 \right)^{p/2} \\ &\leq [2\varepsilon \mathcal{E}_0]^{1/q} \cdot \left(h^2 [4(\kappa^3 + 1) \mathcal{E}_0] \right)^{p/2}. \end{aligned}$$

Therefore,

$$\limsup_{h \rightarrow 0} \frac{1}{h} \left(\int_J \left| u(\tau + h, x) - u(\tau, x) - h \right|^p dx \right)^{1/p} \leq [2\varepsilon \mathcal{E}_0]^{1/pq} \cdot [4(\kappa^3 + 1) \mathcal{E}_0]^{1/2}. \quad (6.8)$$

In a similar way we estimate

$$\begin{aligned} \int_J |u_t(\tau, x)|^p dx &\leq [\text{meas}(J)]^{1/q} \cdot \left(\int_J |u_t(\tau, x)|^2 dx \right)^{p/2}, \\ \left(\int_J |u_t(\tau, x)|^p dx \right)^{1/p} &\leq \text{meas}(J)^{1/pq} \cdot [\kappa^2 \mathcal{E}_0]^{p/2}. \end{aligned} \quad (6.9)$$

Since $\varepsilon > 0$ is arbitrary, from (6.6), (6.8) and (6.9) we conclude

$$\lim_{h \rightarrow 0} \frac{1}{h} \left(\int_{\mathbb{R}} \left| u(\tau + h, x) - u(\tau, x) - h u_t(\tau, x) \right|^p dx \right)^{1/p} = 0. \quad (6.10)$$

The proof of continuity of the map $t \mapsto u_t$ is similar. Fix $\varepsilon > 0$. Consider the intervals $[a_i, b_i]$ as before. Since u is smooth on a neighborhood of $\{\tau\} \times J'$, it suffices to estimate

$$\begin{aligned} & \limsup_{h \rightarrow 0} \int |u_t(\tau + h, x) - u_t(\tau, x)|^p dx \\ & \leq \limsup_{h \rightarrow 0} \int_J |u_t(\tau + h, x) - u_t(\tau, x)|^p dx \\ & \leq \limsup_{h \rightarrow 0} [\text{meas}(J)]^{1/q} \cdot \left(\int_J |u_t(\tau + h, x) - u_t(\tau, x)|^2 dx \right)^{p/2} \\ & \leq \limsup_{h \rightarrow 0} [2\varepsilon \mathcal{E}_0]^{1/q} \cdot \left(\|u_t(\tau + h, \cdot)\|_{\mathbf{L}^2} + \|u_t(\tau, \cdot)\|_{\mathbf{L}^2} \right)^p \\ & \leq [2\varepsilon \mathcal{E}_0]^{1/q} [4\mathcal{E}_0]^p. \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, this proves continuity.

To extend the result to general initial data, such that $(u_0)_x, u_1 \in \mathbf{L}^2$, we consider a sequence of smooth initial data, with $(u_0^n)_x, u_1^n \in \mathcal{C}_c^\infty$, with $u_0^n \rightarrow u_0$ uniformly, $(u_0^n)_x \rightarrow (u_0)_x$ almost everywhere and in \mathbf{L}^2 , $u_1^n \rightarrow u_1$ almost everywhere and in \mathbf{L}^2 .

The continuity of the function $t \mapsto u_x(t, \cdot)$ as a map with values in \mathbf{L}^p , $1 \leq p < 2$, is proved in an entirely similar way.

7 Energy conservation

This section is devoted to the proof of Theorem 3, stating that, in some sense, the total energy of the solution remains constant in time.

A key tool in our analysis is the *wave interaction potential*, defined as

$$\Lambda(t) \doteq (\mu_t^- \otimes \mu_t^+) \{(x, y); x > y\}. \quad (7.1)$$

We recall that μ_t^\pm are the positive measures defined at (5.13). Notice that, if μ_t^+, μ_t^- are absolutely continuous w.r.t. Lebesgue measure, so that (5.10) holds, then (7.1) is equivalent to

$$\Lambda(t) \doteq \frac{1}{4} \iint_{x>y} R^2(t, x) S^2(t, y) dx dy.$$

Lemma 1. *The map $t \mapsto \Lambda(t)$ has locally bounded variation. Indeed, there exists a one-sided Lipschitz constant L_0 such that*

$$\Lambda(t) - \Lambda(s) \leq L_0 \cdot (t - s) \quad t > s > 0. \quad (7.2)$$

To prove the lemma, we first give a formal argument, valid when the solution $u = u(t, x)$ remains smooth. We first notice that (2.4) implies

$$\begin{aligned} \frac{d}{dt}(4\Lambda(t)) &\leq - \int 2c R^2 S^2 dx + \int (R^2 + S^2) dx \cdot \int \frac{c'}{2c} |R^2 S - RS^2| dx \\ &\leq -2\kappa^{-1} \int R^2 S^2 dx + 4\mathcal{E}_0 \left\| \frac{c'}{2c} \right\|_{\mathbf{L}^\infty} \int |R^2 S - RS^2| dx, \end{aligned}$$

where κ^{-1} is a lower bound for $c(u)$. For each $\varepsilon > 0$ we have $|R| \leq \varepsilon^{-1/2} + \varepsilon^{1/2} R^2$. Choosing $\varepsilon > 0$ such that

$$\kappa^{-1} > 4\mathcal{E}_0 \left\| \frac{c'}{2c} \right\|_{\mathbf{L}^\infty} \cdot 2\sqrt{\varepsilon},$$

we thus obtain

$$\frac{d}{dt}(4\Lambda(t)) \leq -\kappa^{-1} \int R^2 S^2 dx + \frac{16 \mathcal{E}_0^2}{\sqrt{\varepsilon}} \left\| \frac{c'}{2c} \right\|_{\mathbf{L}^\infty}.$$

This yields the \mathbf{L}^1 estimate

$$\int_0^\tau \int (|R^2 S| + |RS^2|) dx dt = \mathcal{O}(1) \cdot [\Lambda(0) + \mathcal{E}_0^2 \tau] = \mathcal{O}(1) \cdot (1 + \tau) \mathcal{E}_0^2,$$

where $\mathcal{O}(1)$ denotes a quantity whose absolute value admits a uniform bound, depending only on the function $c = c(u)$ and not on the particular solution under consideration. In particular, the map $t \mapsto \Lambda(t)$ has bounded variation on any bounded interval. It can be discontinuous, with downward jumps.

To achieve a rigorous proof of Lemma 1, we need to reproduce the above argument in terms of the variables X, Y . As a preliminary, we observe that for every $\varepsilon > 0$ there exists a constant κ_ε such that

$$\begin{aligned} &|\sin z(1 - \cos w) - \sin w(1 - \cos z)| \\ &\leq \kappa_\varepsilon \cdot \left(\tan^2 \frac{w}{2} + \tan^2 \frac{z}{2} \right) (1 + \cos w)(1 + \cos z) + \varepsilon(1 - \cos w)(1 - \cos z) \end{aligned} \tag{7.3}$$

for every pair of angles w, z .

Now fix $0 \leq s < t$. Consider the sets Γ_s, Γ_t as in (5.11) and define $\Gamma_{st} \doteq \Gamma_t \setminus \Gamma_s$. Observing that

$$dx dt = \frac{pq}{8c} (1 + \cos w)(1 + \cos z) dX dY,$$

we can now write

$$\begin{aligned} \int_s^t \int_{-\infty}^\infty \frac{R^2 + S^2}{4} dx dt &= (t - s) \mathcal{E}_0 \\ &= \iint_{\Gamma_{st}} \frac{1}{4} \left(\tan^2 \frac{w}{2} + \tan^2 \frac{z}{2} \right) \cdot \frac{pq}{8c} (1 + \cos w)(1 + \cos z) dX dY. \end{aligned} \tag{7.4}$$

The first identity holds only for smooth solutions, but the second one is always valid. Recalling (5.4) and (5.13), and then using (7.3)-(7.4), we obtain

$$\begin{aligned}
\Lambda(t) - \Lambda(s) &\leq - \iint_{\Gamma_{st}} \frac{1-\cos w}{8} p \cdot \frac{1-\cos z}{8} q \, dX dY \\
&\quad + \mathcal{E}_0 \cdot \iint_{\Gamma_{st}} \frac{c'}{64c^2} pq [\sin z(1 - \cos w) - \sin w(1 - \cos z)] \, dX dY \\
&\leq -\frac{1}{64} \iint_{\Gamma_{st}} (1 - \cos w)(1 - \cos z) pq \, dX dY \\
&\quad + \mathcal{E}_0 \cdot \iint_{\Gamma_{st}} \frac{c'}{64c^2} pq \left[\kappa_\varepsilon \cdot \left(\tan^2 \frac{w}{2} + \tan^2 \frac{z}{2} \right) (1 + \cos w)(1 + \cos z) \right. \\
&\quad \left. + \varepsilon(1 - \cos w)(1 - \cos z) \right] \, dX dY \\
&\leq \kappa(t - s),
\end{aligned}$$

for a suitable constant κ . This proves the lemma.

To prove Theorem 3, consider the three sets

$$\begin{aligned}
\Omega_1 &\doteq \left\{ (X, Y); \quad w(X, Y) = -\pi, \quad z(X, Y) \neq -\pi, \quad c'(u(X, Y)) \neq 0 \right\}, \\
\Omega_2 &\doteq \left\{ (X, Y); \quad z(X, Y) = -\pi, \quad w(X, Y) \neq -\pi, \quad c'(u(X, Y)) \neq 0 \right\}, \\
\Omega_3 &\doteq \left\{ (X, Y); \quad z(X, Y) = -\pi, \quad w(X, Y) = -\pi, \quad c'(u(X, Y)) \neq 0 \right\}.
\end{aligned}$$

From the equations (2.24), it follows that

$$\text{meas}(\Omega_1) = \text{meas}(\Omega_2) = 0. \quad (7.5)$$

Indeed, $w_Y \neq 0$ on Ω_1 and $z_X \neq 0$ on Ω_2 .

Let Ω_3^* be the set of Lebesgue points of Ω_3 . We now show that

$$\text{meas} \left(\{t(X, Y); (X, Y) \in \Omega_3^*\} \right) = 0. \quad (7.6)$$

To prove (7.4), fix any $P^* = (X^*, Y^*) \in \Omega_3^*$ and let $\tau = t(P^*)$. We claim that

$$\limsup_{h, k \rightarrow 0^+} \frac{\Lambda(\tau - h) - \Lambda(\tau + k)}{h + k} = +\infty \quad (7.7)$$

By assumption, for any $\varepsilon > 0$ arbitrarily small we can find $\delta > 0$ with the following property. For any square Q centered at P^* with side of length $\ell < \delta$, there exists a vertical segment σ and a horizontal segment σ' , as in fig. 6, such that

$$\text{meas}(\Omega_3 \cap \sigma) \geq (1 - \varepsilon)\ell, \quad \text{meas}(\Omega_3 \cap \sigma') \geq (1 - \varepsilon)\ell, \quad (7.8)$$

Call

$$t^+ \doteq \max \left\{ t(X, Y); (X, Y) \in \sigma \cup \sigma' \right\},$$




Figure 6:

$$t^- \doteq \min \left\{ t(X, Y); (X, Y) \in \sigma \cup \sigma' \right\}.$$

Notice that, by (4.2),

$$t^+ - t^- \leq \int_{\sigma} \frac{(1 + \cos w)p}{4c} dX + \int_{\sigma'} \frac{(1 + \cos z)q}{4c} dY \leq c_0 \cdot (\varepsilon \ell)^2. \quad (7.9)$$

Indeed, the integrand functions are Lipschitz continuous. Moreover, they vanish outside a set of measure $\varepsilon \ell$. On the other hand,

$$\Lambda(t^-) - \Lambda(t^+) \geq c_1(1 - \varepsilon)^2 \ell^2 - c_2(t^+ - t^-) \quad (7.10)$$

for some constant $c_1 > 0$. Since $\varepsilon > 0$ was arbitrary, this implies (7.5).

Recalling that the map $t \mapsto \Lambda$ has bounded variation, from (7.5) it follows (7.4).

We now observe that the singular part of μ_τ is nontrivial only if the set

$$\{P \in \gamma_\tau; w(P) = -\pi \text{ or } z(P) = -\pi\}$$

has positive 1-dimensional measure. By the previous analysis, restricted to the region where $c' \neq 0$, this can happen only for a set of times having zero measure.

Acknowledgment: Alberto Bressan was supported by the Italian M.I.U.R., within the research project #2002017219, while Yuxi Zheng has been partially supported by grants NSF DMS 0305497 and 0305114.

References

- [1] M. Albers, R. Camassa, D. Holm, and J. Marsden, The geometry of peaked solitons and billiard solutions of a class of integrable PDE's, *Lett. Math. Phys.*, **32**(1994), pp. 137–151.

- [2] M. Balabane, Non-existence of global solutions for some nonlinear wave equations with small Cauchy data, *C. R. Acad. Sc. Paris*, **301**(1985), pp. 569–572.
- [3] H. Berestycki, J. M. Coron and I. Ekeland (eds.), *Variational Methods*, Progress in Nonlinear Differential Equations and Their Applications, Vol. 4, Birkhäuser, Boston (1990).
- [4] A. Bressan and A. Constantin, Global solutions to the Camassa-Holm equations, to appear.
- [5] A. Bressan, Ping Zhang, and Yuxi Zheng, On asymptotic variational wave equations, *Arch. Rat. Mech. Anal.*, submitted April 29, 2004.
- [6] R. Camassa and D. Holm, An integrable shallow water equation with peaked solitons, *Phys. Rev. Lett.*, **71**(1993), pp. 1661–1664.
- [7] R. Camassa, D. Holm, J. Hyman, A new integrable shallow water equation, to appear in *Adv. in Appl. Mech.*
- [8] D. Christodoulou and A. Tahvildar-Zadeh, On the regularity of spherically symmetric wave maps, *Comm. Pure Appl. Math.*, **46**(1993), pp. 1041–1091.
- [9] J. Coron, J. Ghidaglia, and F. Hélein (eds.), *Nematics*, Kluwer Academic Publishers, 1991.
- [10] J. L. Ericksen and D. Kinderlehrer (eds.), *Theory and Application of Liquid Crystals*, IMA Volumes in Mathematics and its Applications, Vol. 5, Springer-Verlag, New York (1987).
- [11] R. T. Glassey, Finite-time blow-up for solutions of nonlinear wave equations, *Math. Z.*, **177**(1981), pp. 323–340.
- [12] R. T. Glassey, J. K. Hunter and Yuxi Zheng, Singularities in a nonlinear variational wave equation, *J. Differential Equations*, **129**(1996), 49–78.
- [13] R. T. Glassey, J. K. Hunter and Yuxi Zheng, Singularities and oscillations in a nonlinear variational wave equation, *Singularities and Oscillations*, edited by J. Rauch and M. E. Taylor, IMA, Vol **91**, Springer, 1997.
- [14] A. Grundland and E. Infeld, A family of nonlinear Klein-Gordon equations and their solutions, *J. Math. Phys.*, **33**(1992), pp. 2498–2503.
- [15] B. Hanouzet and J. L. Joly, Explosion pour des problèmes hyperboliques semi-linéaires avec second membre non compatible, *C. R. Acad. Sc. Paris*, **301**(1985), pp. 581–584.
- [16] R. Hardt, D. Kinderlehrer, and Fanghua Lin, Existence and partial regularity of static liquid crystal configurations. *Comm. Math. Phys.*, **105**(1986), pp. 547–570.

- [17] J. K. Hunter and R. A. Saxton, Dynamics of director fields, *SIAM J. Appl. Math.*, **51**(1991), pp. 1498-1521.
- [18] J. K. Hunter and Yuxi Zheng, On a nonlinear hyperbolic variational equation I and II, *Arch. Rat. Mech. Anal.*, **129**(1995), pp. 305-353 and 355-383.
- [19] J. K. Hunter and Yuxi Zheng, On a completely integrable nonlinear hyperbolic variational equation, *Physica D*. **79**(1994), 361–386.
- [20] F. John, Blow-up of solutions of nonlinear wave equations in three space dimensions, *Manuscripta Math.*, **28**(1979), pp. 235–268.
- [21] T. Kato, Blow-up of solutions of some nonlinear hyperbolic equations, *Comm. Pure Appl. Math.*, **33**(1980), pp. 501–505.
- [22] D. Kinderlehrer, Recent developments in liquid crystal theory, in *Frontiers in pure and applied mathematics : a collection of papers dedicated to Jacques-Louis Lions on the occasion of his sixtieth birthday*, ed. R. Dautray, Elsevier, New York, pp. 151–178 (1991).
- [23] S. Klainerman and M. Machedon, Estimates for the null forms and the spaces $H_{s,\delta}$, *Internat. Math. Res. Notices*, 1996, no. 17, pp. 853–865.
- [24] S. Klainerman and A. Majda, Formation of singularities for wave equations including the nonlinear vibrating string, *Comm. Pure Appl. Math.*, **33**(1980), pp. 241–263.
- [25] P. Lax, Development of singularities of solutions of nonlinear hyperbolic partial differential equations, *J. Math. Phys.*, **5**(1964), pp. 611–613.
- [26] H. Levine, Instability and non-existence of global solutions to nonlinear wave equations, *Trans. Amer. Math. Soc.*, **192**(1974), pp. 1–21.
- [27] H. Lindblad, Global solutions of nonlinear wave equations, *Comm. Pure Appl. Math.*, **45**(1992), pp. 1063–1096.
- [28] Tai-Ping Liu, Development of singularities in the nonlinear waves for quasi-linear hyperbolic partial differential equations, *J. Differential Equations*, **33**(1979), pp. 92–111.
- [29] R. A. Saxton, Dynamic instability of the liquid crystal director, in *Contemporary Mathematics* Vol. 100: Current Progress in Hyperbolic Systems, pp. 325–330, ed. W. B. Lindquist, AMS, Providence, 1989.
- [30] J. Schaeffer, The equation $u_{tt} - \Delta u = |u|^p$ for the critical value of p , *Proc. Roy. Soc. Edinburgh Sect. A*, **101A**(1985), pp. 31–44.

- [31] J. Shatah, Weak solutions and development of singularities in the $SU(2)$ σ -model, *Comm. Pure Appl. Math.*, **41**(1988), pp. 459–469.
- [32] J. Shatah and A. Tahvildar-Zadeh, Regularity of harmonic maps from Minkowski space into rotationally symmetric manifolds, *Comm. Pure Appl. Math.*, **45**(1992), pp. 947–971.
- [33] T. Sideris, Global behavior of solutions to nonlinear wave equations in three dimensions, *Comm. in Partial Diff. Eq.*, **8**(1983), pp. 1291–1323.
- [34] T. Sideris, Nonexistence of global solutions to semilinear wave equations in high dimensions, *J. Diff. Eq.*, **52**(1984), pp. 378–406.
- [35] W. Strauss, *Nonlinear wave equations*, CBMS Lectures 73, AMS, Providence, 1989.
- [36] E. Virga, *Variational Theories for Liquid Crystals*, Chapman & Hall, New York (1994).
- [37] Ping Zhang and Yuxi Zheng, On oscillations of an asymptotic equation of a nonlinear variational wave equation, *Asymptotic Analysis*, **18**(1998), pp. 307–327 .
- [38] Ping Zhang and Yuxi Zheng, On the existence and uniqueness of solutions to an asymptotic equation of a variational wave equation, *Acta Mathematica Sinica*, **15**(1999), pp. 115–130.
- [39] Ping Zhang and Yuxi Zheng, On the existence and uniqueness to an asymptotic equation of a variational wave equation with general data, *Arch. Rat. Mech. Anal.* **155**(2000), 49–83.
- [40] Ping Zhang and Yuxi Zheng, Rarefactive solutions to a nonlinear variational wave equation, *Comm. Partial Differential Equations*, **26**(2001), 381–419.
- [41] Ping Zhang and Yuxi Zheng, Singular and rarefactive solutions to a nonlinear variational wave equation, *Chinese Annals of Mathematics*, **22B**, 2(2001), 159–170.
- [42] Ping Zhang and Yuxi Zheng, Weak solutions to a nonlinear variational wave equation, *Arch. Rat. Mech. Anal.*, **166** (2003), 303–319.
- [43] Ping Zhang and Yuxi Zheng, On the second-order asymptotic equation of a variational wave equation, *Proc A of the Royal Soc. Edinburgh, A. Mathematics*, **132A**(2002), 483–509.
- [44] Ping Zhang and Yuxi Zheng, Weak solutions to a nonlinear variational wave equation with general data, *Annals of Inst. H. Poincaré*, 2004 (in press).
- [45] H. Zorski and E. Infeld, New soliton equations for dipole chains, *Phys. Rev. Lett.*, **68**(1992), pp. 1180–1183.